

Lactuca chondrilliflorae-Artemisietum campestris Billy ex Loiseau & Felzines 2010

																				SYNTHESE
	Numéro relevé	222832	222834	222835	222836	222927	222928	222930	222937	222938	222941	386057	386059	386071	386092	386094	61574			
	Altitude (m)	420	400	700	450	650	480	580	800	660	500	470	810	880	870	720	620		626	
	Pente relevé (°)	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd		nd	
	Exposition																			
	Surface (m²)	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	20		20	
	Recouvrement total (%)	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	90		90	
	Nombre de taxons	10	13	16	18	17	20	15	21	16	20	21	35	30	14	25	48		21	
Combinaison caractéristique	<i>Artemisia campestris</i> L.	.	.	1	1,2	1	1	2	1	2	1	1	1,2	1,2	2,3	2	1		V ¹⁻²	
	<i>Lactuca viminea</i> (L.) J.Presl & C.Presl subsp. <i>chondrilliflora</i> (Boreau) Bonnier	+	+	1	1	1	1	1,2	+	1	1	1	+	1	1	1,2	+		V ⁺¹	
	<i>Sedum album</i> L.	1	1,2	1,2	1,2	2	2	2	.	2	1	2	1	1,2	.	1	1		V ¹⁻²	
	<i>Centaurea stoebe</i> L.	+	+	.	1	1	1	1	.	.	+	+	+	.	.	.	1		III ⁺¹	
	<i>Festuca arvensensis</i> Auquier, Kerguelén & Markgr.-Dann.	.	.	1	1	1,2	.	.	2,3	1,2	1,2	.	3	1,2	1	2	1		III ¹⁻³	
	<i>Sedum rupestre</i> L.	.	.	1	.	1	.	.	.	.	.	1	1	1	.	1	1		III ¹	
	<i>Stachys recta</i> L.	.	.	.	1	+	.	1	.	1	1	.	1	.	.	.	1		III ⁺¹	
	<i>Melica ciliata</i> L.	.	.	.	1	+	1	1	1	1,2	1	1	.	.	.	.	1		III ⁺¹	
	<i>Oreoselinum nigrum</i> Delarbre	+	1	.	.	1	.	.	2	.	.	.	.	.	.	2	.		II ⁺²	
	<i>Echium vulgare</i> L.	.	.	.	+	.	1	+	.	.	.	1	+	+	.	.	1		III ⁺¹	
<i>Andryala integrifolia</i> L.	.	.	.	1	.	1	1	.	.	.	1	.	.	.	.	1		II ¹		
<i>Anthemis cretica</i> L.	1,2	1,2	1,2	1,2	.	.	.	.	.	.	.	.	.	.	.	.	.		II ¹	
Espèces des milieux secs très maigres	<i>Thymus gr. praecox</i>	.	1	1	+	1,2	1	1	1	1	1,2	1	1,2	1	.	.	2		IV ⁺²	
	<i>Hylotelephium maximum</i> (L.) Holub	1	+	.	.	.	.	.	.	.	.	.	.	.	.	.	.		I ⁺¹	
	<i>Fumana procumbens</i> (Dunal) Gren. & Godr.	.	.	.	.	.	+	.	.	.	.	+	.	.	.	.	.		I ⁺	
	<i>Hippocrepis comosa</i> L.	.	.	.	.	.	.	.	.	.	1	.	.	.	+	.	1		I ⁺¹	
	<i>Koeleria vallesiana</i> (Honck.) Gaudin	.	.	.	.	.	.	.	.	1	1	.	.	+	.	.	.		I ⁺¹	
Espèces des milieux secs, très maigres à maigres	<i>Sedum acre</i> L.	.	.	.	.	.	.	.	.	1	.	.	.	.	.	.	.		I ¹	
	<i>Potentilla gr. argentea</i>	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		+	
	<i>Jasione montana</i> L.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	+		I ⁺	
	<i>Phleum phleoides</i> (L.) H.Karst.	.	.	.	.	.	.	.	1	.	1,2	.	1,2	.	+	1	2		II ⁺²	
	<i>Allium sphaerocephalon</i> L.	.	.	.	.	+	2	.	.	+	.	2	.	.	+	.	2		II ⁺²	
	<i>Trifolium striatum</i> L.	.	.	.	.	.	.	.	.	.	.	.	.	1	.	.	1		I ¹	
	<i>Anarrhinum bellidifolium</i> (L.) Willd.	+	+	1	+	.	+	.	.	.	.	.	.	.	.	.	.		II ⁺¹	
	<i>Koeleria gr. pyramidata</i>	.	.	.	.	.	.	.	1	.	.	.	1	.	.	1	1		I ¹	
	Espèces des milieux secs à sains, très maigres à maigres	R- <i>Anthyllis vulneraria</i> L.	.	.	.	.	.	.	.	.	.	.	.	1	.	.	.	.		I ¹
<i>Poterium sanguisorba</i> L.		.	.	+	1	+	+	.	.	1	1	+	1	+	.	+	+		IV ⁺¹	
<i>Potentilla verna</i> L.		.	.	+	.	.	1	1	1	.	1	1	1,2	1	1	1	1		IV ⁺¹	
<i>Helianthemum nummularium</i> (L.) Mill.		.	.	.	.	.	1	.	1	.	1	1	1,2	1	+	1	1		III ⁺¹	
<i>Rumex acetosella</i> L.		.	.	.	+	.	.	+	.	.	.	.	.	.	.	.	.		I ⁺	
<i>Genista sagittalis</i> L.		.	.	.	.	.	.	+	+	.	.	.	.	.	.	.	.		I ⁺	
Ci <i>Galium album</i> Mill.		.	.	.	.	.	.	.	.	.	.	.	.	.	1	.	1		I ¹	
<i>Campanula rotundifolia</i> L.		.	.	.	.	.	.	.	+	.	.	.	1	.	.	.	.		I ⁺¹	
<i>Inula montana</i> L.		.	.	.	.	.	.	.	.	+	.	.	.	.	.	.	.		I ⁺	
<i>Silene nutans</i> L.		.	.	.	.	1	.	1	.	.	.	.	.	.	.	1	.		I ¹	
Espèces des milieux secs à sains, très maigres à peu fertiles	<i>Euphorbia cyparissias</i> L.	+	+	.	+	.	+	1	1	.	1	+	1	+	.	1	1		IV ⁺¹	
	<i>Hypericum perforatum</i> L.	.	.	.	.	.	.	.	+	.	.	.	1	1	+	1	1		II ⁺¹	
	<i>Eryngium campestre</i> L.	.	.	.	.	.	.	.	+	+	1	.	1	+	+	1	1		III ⁺¹	
	<i>Dianthus carthusianorum</i> L.	.	.	.	.	.	.	.	.	.	+	.	1	+	.	1	+		II ⁺¹	
	<i>Poa bulbosa</i> L.	.	.	.	.	.	.	.	.	+	.	.	.	1	.	.	.		I ⁺¹	
	<i>Scabiosa columbaria</i> L.	.	.	.	+	.	+	.	.	.	+	.	1	.	.	+	1		II ⁺¹	
	Espèces des milieux secs à sains, maigres à peu fertiles	<i>Chondrilla juncea</i> L.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		+
<i>Galium verum</i> L.		.	.	.	.	.	.	.	.	.	.	.	1	.	.	.	.		I ¹	
<i>Rhinanthus minor</i> L.		.	.	.	.	.	.	.	.	.	.	.	1	.	.	.	.		I ¹	
R- <i>Brachypodium rupestre</i> (Host) Roem. & Schult.		.	.	.	.	.	.	.	.	.	.	.	.	.	.	1	.		I ¹	
<i>Saxifraga granulata</i> L.		.	.	.	.	.	.	.	.	.	.	.	.	1	.	.	.		I ¹	
<i>Teucrium scorodonia</i> L.		.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	1		I ¹	
<i>Asperula cynanchica</i> L.		.	.	.	.	.	.	.	.	.	.	.	.	+	.	.	.		I ⁺	
<i>Ononis spinosa</i> L.		.	.	.	.	.	.	.	+	.	.	.	.	.	.	.	.		I ⁺	
C <i>Odontites vernus</i> (Bellardi) Dumort.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	+		I ⁺		
Espèces à large amplitude	<i>Achillea millefolium</i> L.	.	.	.	.	.	.	.	1	.	.	.	1	+	1	1	1		II ⁺¹	
	<i>Plantago lanceolata</i> L.	.	.	.	.	.	.	.	.	.	.	.	.	+	.	.	.		I ⁺	
	<i>Agrostis capillaris</i> L.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	1		I ¹	
Espèces annuelles	<i>Trifolium arvense</i> L.	.	.	.	.	+	.	+	+	.	.	.	1	1,2	.	+	1		III ⁺¹	
	<i>Trifolium campestre</i> Schreb.	.	.	.	.	.	.	.	1	.	.	1	1	+	.	1	.		II ⁺¹	
Autres taxons	F <i>Arrhenatherum elatius</i> (L.) P.Beauv. ex J.Presl & C.Presl	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	1		I ¹	
	<i>Teucrium chamaedrys</i> L.	.	.	.	.	.	.	.	.	.	.	.	.	.	2	.	.		I ²	
	<i>Lotus corniculatus</i> L.	.	.	.	.	.	.	.	.	.	.	.	+	.	.	.	.		I ⁺	
	<i>Poa bulbosa</i> L.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	1		I ¹	
	<i>Vicia lutea</i> L.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	1		I ¹	
		.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.			

	Numéro relevé	222832	222834	222835	222836	222927	222928	222930	222937	222938	222941	386057	386059	386071	386092	386094	815574	626	SYNTHESE								
	Altitude (m)	420	400	700	450	650	480	580	800	660	500	470	810	880	870	720	620	20	20								
	Pente relevé (°)	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	90	90								
	Exposition																										
	Surface (m²)	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	20	20									
	Recouvrement total (%)	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	90	90									
	Nombre de taxons	10	13	16	18	17	20	15	21	16	20	21	35	30	14	25	48	21	21								
Ranunculus bulbosus L.		.	.	.	.	.	.	.	.	.	.	.	.	1	.	.	+	I ⁺¹									
Muscari comosum (L.) Mill.		.	.	.	.	+	+	.	.	.	.	+	.	.	.	.	.	I ⁺									
Sempervivum tectorum L.		.	.	.	.	.	.	.	.	1	.	.	.	.	.	.	.	I ¹									
Lactuca perennis L.		.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	1	I ¹									
Clinopodium acinos (L.) Kuntze		.	.	.	.	.	.	.	+	.	.	.	.	.	.	.	.	I ⁺									
Silene otites (L.) Wibel		.	.	.	.	.	.	.	.	.	.	.	1	.	.	.	.	I ¹									
Atocion armeria (L.) Raf.		.	+	.	1	.	.	.	.	.	.	.	.	.	.	.	.	I ⁺¹									
Bromus squarrosus L.		.	.	.	.	.	.	.	.	.	.	+	.	.	.	.	.	I ⁺									
Asplenium septentrionale (L.) Hoffm.		1	1	+	.	.	.	.	.	.	.	.	.	.	.	.	.	I ⁺¹									
Crucianella angustifolia L.		.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	1	I ¹									
Sedum hirsutum All.		.	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	I ¹									
Orobanche amethystea Thuill.		.	.	.	.	.	.	.	.	.	.	.	.	.	.	+	.	I ⁺									
Hylotelephium telephium (L.) H.Ohba		.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	I ⁺									
Saponaria ocymoides L.		.	.	.	.	1,2	.	.	.	.	.	.	.	.	.	.	.	I ⁺									
Allium flavum L.		.	.	.	.	.	.	.	.	.	.	.	1	.	.	.	.	I ¹									
Cistus umbellatus L.		.	.	1	.	.	.	.	.	.	.	.	.	.	.	.	.	I ¹									
Rubus ulmifolius Schott		.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	2	I ²									
Orobanche alba Stephan ex Willd.		.	.	.	.	.	.	.	.	.	.	+	.	.	.	.	.	I ⁺									
Arenaria serpyllifolia L.		.	.	.	.	.	.	.	.	.	.	.	+	+	.	.	.	I ⁺									
Logfia minima (Sm.) Dumort.		.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	1	I ¹									
Medicago minima (L.) L.		.	.	.	.	.	.	.	.	.	.	.	1	.	.	.	.	I ¹									
Trifolium glomeratum L.		.	.	.	.	.	.	.	.	.	.	.	.	+	.	.	.	I ⁺									
Aira caryophyllea L.		.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	1	I ¹									
Alyssum alyssoides (L.) L.		.	.	.	.	.	.	.	.	.	.	.	1	.	.	.	.	I ¹									
Ervilia hirsuta (L.) Opiz		.	.	.	.	.	.	.	.	.	.	.	.	1	.	.	.	I ¹									
Petrorhagia prolifera (L.) P.W.Ball & Heywood		.	.	.	.	.	.	.	.	.	.	.	.	+	.	.	.	I ⁺									
Micropyrum tenellum (L.) Link		.	.	+	.	.	+	.	.	.	.	.	.	.	.	.	1	I ⁺¹									
Veronica arvensis L.		.	.	.	.	.	.	.	.	.	.	+	+	.	.	.	.	I ⁺									
Draba verna L.		.	.	.	.	.	.	.	.	.	.	.	.	+	.	.	.	I ⁺									
Cerastium pumilum Curtis		.	.	+	.	.	.	.	.	.	.	.	+	.	.	.	.	I ⁺									
Teesdalia nudicaulis (L.) R.Br.		.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	1	I ¹									
Genista pilosa L.		.	.	.	.	.	.	.	.	.	.	.	.	.	1	.	.	I ¹									
Cytisus oromediterraneus Rivas Mart. & al.		1	1	1	.	.	.	.	.	.	.	.	.	.	.	.	2	II ¹⁻²									
Erigeron canadensis L.		.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	+	I ⁺									
Cytisus scoparius (L.) Link		.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	+	I ⁺									
Prunus spinosa L.		.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	2	I ²									