

Teucrio chamaedrys-Brometum erecti

Billy ex Thébaud, C. Roux, C.-E. Bernard & Delcoigne 2014

		SYNTHÈSE																								
		Numéro relevé				222488	222489	222491	222492	222494	333335	333337	333338	333355	333358	333364	333369	333376	333378	333380	333385	333389	333395	SYNTHÈSE		
		Altitude (m)				500	520	360	460	630	700	640	447	380	400	420	460	360	610	460	620	560	480	500		
		Pente relevé (°)				nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd		
		Exposition																								
		Surface (m²)				nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd		
		Recouvrement total (%)				nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd		
		Nombre de taxons				22	26	20	28	32	16	27	22	31	38	36	25	24	29	28	37	23	23	27		
Combinaison caractéristique																										
R-	<i>Bromopsis erecta</i> (Huds.) Fourr.	1	3	1	2	4	.	1	2	3	1,2	2	1	1,2	3	2	3	1	4						V ⁺¹⁻⁴	
	<i>Brachypodium rupestre</i> (Host) Roem. & Schult.	.	1	+	2	1	.	1	1	3	4	1,2	1	3	4	2,3	3	1	1						V ⁺⁴	
	<i>Helianthemum apenninum</i> (L.) Mill.	1,2	1	1	1	+	1,2	.	.	+	+	1	.	1	+	+	1	.							IV ⁺¹	
	<i>Teucrium chamaedrys</i> L.	1	+	1	+	1	2	1,2	1,2	.	.	+	.	+	.	.	.	.	1,2	+					IV ⁺²	
	<i>Hippocrepis comosa</i> L.	2	+	2	2	2	.	.	2	+	+	1	.	.	1,2	.	3	2	1						IV ⁺³	
	<i>Thymus gr. praecox</i>	+	+	.	.	+	+	1	1,2	+	+	.	1	.	1	.	+	+							IV ⁺¹	
	<i>Festuca lemanii</i> Bastard	1 cf	1 cf	1 cf	.	.	1	1,2	1	+	+	1,2	.	.	1,2	.	.	.	1,2	.						IV ⁺¹
	<i>Galium glaucum</i> L.	1,2	.	.	+	.	.	1	.	.	1,2	1	+	+	.	.	.	.	1	.						III ⁺¹
	<i>Onobrychis viciifolia</i> Scop.	.	1	+	1	1	.	.	1	.	+	.	.	.	.	.	+	1	1	1	+					III ⁺¹
	<i>Koeleria vallesiana</i> (Honck.) Gaudin	.	.	.	.	.	.	1	.	.	+	.	.	.	.	.	.	.	.	+	1					II ⁺¹
C	<i>Inula bifrons</i> (L.) L.	.	.	.	.	.	.	.	.	+	+	+	.	+	.	.	.	.	.	.	.	.	.	.		II ⁺
	<i>Linum tenuifolium</i> L.	.	.	+	.	+	.	.	+	.	.	.	.	.	.	.	+	.	.	.	.	.	.	.		II ⁺
	<i>Plantago media</i> L.	.	.	.	.	.	.	.	.	.	.	.	+	.	+	+	+	+	.	.	.	.	.	.		II ⁺
	<i>Fumana procumbens</i> (Dunal) Gren. & Godr.	.	.	.	.	.	1	.	1	.	.	.	.	.	.	.	.	.	+	.	.	.	.	.		I ⁺¹
	Espèces des milieux secs, très maigres à maigres																									
	<i>Stachys recta</i> L.	+	+	+	.	.	1	+	.	.	.	.	.	.	.	.	.	.	.	.	1	.	.	.		II ⁺¹
	<i>Coronilla minima</i> L.	.	.	2	.	1	.	.	.	.	.	.	.	.	.	.	.	1	.	.	.	.	.	.		I ¹⁻²
	<i>Phleum phleoides</i> (L.) H.Karst.	.	.	.	.	.	1	.	.	.	.	.	.	.	.	.	.	+	.	.	.	.	.	.		I ⁺¹
	Espèces des milieux secs à sains, très maigres à maigres																									
	Ci	<i>Coronilla varia</i> L.	.	.	.	.	.	.	.	.	.	.	.	.	.	1	.	.	.	.	.	.	.	.	+	
<i>Poterium sanguisorba</i> L.		.	1	+	1	1	.	+	+	+	+	1	+	1	+	1	+	1	1	1	+	.	.			V ⁺¹
<i>Potentilla verna</i> L.		1,2	1	.	+	.	.	1,2	1,2	.	.	1	.	.	1	.	.	1	.	.	1	.	.			III ⁺¹
<i>Trinia glauca</i> (L.) Dumort.		.	.	.	.	.	.	.	.	.	.	+	.	.	.	.	.	.	.	+	+	.	.			I ⁺
<i>Helianthemum nummularium</i> (L.) Mill.		.	.	.	.	.	.	.	1,2	.	.	.	.	.	.	.	.	.	.	.	.	.	.			I ¹
<i>Galium album</i> Mill.		1	.	+	.	.	.	+	.	.	.	.	.	.	.	.	.	.	.	+	+	.	+	.		II ⁺¹
R- <i>Anthyllis vulneraria</i> L.		.	.	.	.	.	.	.	.	.	.	.	.	1,2	.	.	.	.	.	.	1	.	.			I ¹
<i>Carex halleriana</i> Asso		.	.	.	.	.	.	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.			I ¹
<i>Thesium humifusum</i> DC.		.	.	.	.	.	.	.	1	.	.	.	.	.	1	.	+	.	.	.	.	.	.			I ⁺¹
C <i>Ophrys apifera</i> Huds.		.	.	.	.	+	.	.	.	+	+	.	.	.	.	.	.	.	.	.	+	.	.			II ⁺
Espèces des milieux secs à sains, très maigres à peu fertiles																										
C	<i>Hypericum perforatum</i> L.	.	.	.	.	.	.	+	.	.	+	+	.	.	.	.	.	.	.	.	.	.	.			I ⁺
	<i>Cirsium acaulon</i> (L.) Scop.	.	.	.	.	1	.	.	.	.	.	1	1	.	.	1	1	+	.	1	1	+	.			II ⁺¹
	<i>Euphorbia cyparissias</i> L.	1	+	.	+	1	1	+	.	.	+	1	+	1	+	1	+	1	1	1	+	1	.			IV ⁺¹
	<i>Dianthus carthusianorum</i> L.	.	.	+	.	.	.	.	.	.	+	.	.	.	.	.	.	.	.	.	.	.	.			I ⁺
	<i>Ranunculus bulbosus</i> L.	.	+	.	1	1	.	1	.	+	.	1	.	1	.	1	.	.	.	+	+	.	.			III ⁺¹
	<i>Lotus corniculatus</i> L.	+	1	.	.	.	.	+	.	.	+	.	1,2	.	1	.	.	.	.	.	.	.	.			II ⁺¹
	<i>Eryngium campestre</i> L.	+	+	+	+	1,2	+	+	+	+	1	1,2	+	+	+	.	.	.	.	2	.	.	.			IV ⁺²
	<i>Origanum vulgare</i> L.	.	.	.	.	.	.	.	.	.	1	1	.	1	.	1	.	.	.	+	.	.	.			II ⁺¹
	<i>Pilosella officinarum</i> Vaill.	1,2	.	.	1	.	1	.	.	1	1	1	1	1	.	1	1	1	.	.	.	.				III ¹
	<i>Orchis purpurea</i> Huds.	.	.	.	+	+	.	.	.	.	.	.	.	.	.	+	1	1	.	.	.	.	.			II ⁺¹
<i>Poa bulbosa</i> L.	+	.	+	.	.	.	.	.	1	.	.	.	.	.	.	.	.	.	.	.	.	.				I ⁺¹
<i>Scabiosa columbaria</i> L.	.	.	.	.	.	.	.	+	.	.	.	.	.	.	.	.	+	.	.	.	1	.	.			I ⁺¹
Espèces des milieux secs à sains, maigres à peu fertiles																										
Ci	<i>Salvia pratensis</i> L.	.	1,2	.	+	.	1,2	.	.	.	+	.	.	.	.	.	.	.	.	1	.	+	.			II ⁺¹
	<i>Prunella laciniata</i> (L.) L.	.	.	.	.	1	.	.	.	.	.	.	.	.	.	.	.	.	.	1	.	.	.			I ¹
Espèces des milieux secs à frais, très maigres à moyennement fertiles																										
C	<i>Asperula cynanchica</i> L.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	+	.	.	+	.	.	.	.			I ⁺
	<i>Briza media</i> L.	.	.	.	.	1	.	.	.	.	.	.	.	.	+	.	.	+	+	1	.	.	.			II ⁺¹
	<i>Vicia angustifolia</i> L.	.	1	.	+	.	.	.	.	.	.	.	.	.	.	.	1	.	.	.	.	+	.			II ⁺¹
Espèces des milieux sains, très maigres à peu fertiles																										
C	<i>Carlina vulgaris</i> L.	.	.	.	+	+	.	.	.	+	1	+	+	+	+	1	+	1	+	.	.	.	.			III ⁺¹
	<i>Ononis spinosa</i> L.	+	.	.	+	1,2	1	.	+	.	+	+	.	.	+	+	+	1	1	1	1	.	.			IV ⁺¹
	<i>Agrimonia eupatoria</i> L.	.	.	.	1	.	.	.	.	1	1	+	+	+	.	+	+	.	.	.	.	.	.			III ⁺¹
	<i>Carex flacca</i> Schreb.	.	.	.	.	1	.	.	.	.	.	.	.	1	.	.	1	1	1	.	.	+	.			II ⁺¹
	<i>Himantoglossum hircinum</i> (L.) Spreng.	.	+	+	.	+	.	.	.	.	.	+	.	.	.	.	.	.	.	.	+	.	.			II ⁺
C	<i>Blackstonia perfoliata</i> (L.) Huds.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	1	+	.	.	.			I ⁺¹
	<i>Linum catharticum</i> L.	.	.	.	.	+	.	.	.	.	.	.	.	.	.	.	.	.	1	.	.	.	.			I ⁺¹
	<i>Elytrigia intermedia</i> (Host) Nevski	.	.	.	.	.	1	.	.	1	.	1	.	1	.	.	.	.	.	.	.	.	.			I ¹
	<i>Centaurea jacea</i> L.	.	.	.	.	+	.	.	.	+	.	.	+	.	+	.	.	.	1	.	.	.	.			II ⁺¹

			SYNTHESE																		
		Número relevé	222488	222489	222491	222492	222494	383835	383837	383838	383855	383858	383864	383869	383876	383878	383880	383885	383889	383895	500
		Altitude (m)	500	520	360	460	630	700	640	447	380	400	420	460	360	610	460	620	560	480	500
		Pente relevé (°)	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd
		Exposition																			
		Surface (m²)	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd
		Recouvrement total (%)	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd
		Nombre de taxons	22	26	20	28	32	16	27	22	31	38	36	25	24	29	28	37	23	23	27
C	F	<i>Knaulia arvensis</i> (L.) Coult.	.	.	.	.	.	+	.	.	.	.	.	+	.	.	1	.	.	.	I ⁺¹
Espèces à large amplitude																					
		<i>Poa pratensis</i> L.	+	1	+	+	.	.	1	.	1,2	1	.	.	1	.	.	.	.	1	III ⁺¹
		<i>Daucus carota</i> L.	.	1	.	.	+	.	.	.	.	+	.	+	.	.	1	.	.	.	II ⁺¹
Espèces annuelles																					
		<i>Medicago minima</i> (L.) L.	+	+	.	.	.	.	+	.	+	.	.	.	.	.	.	.	.	.	II ⁺
		<i>Arenaria serpyllifolia</i> L.	.	.	.	.	.	.	+	+	+	+	.	.	.	.	.	.	.	.	II ⁺
		<i>Medicago lupulina</i> L.	.	.	.	.	.	.	+	.	+	+	1	.	1	.	.	.	.	+	II ⁺¹
		<i>Trifolium campestre</i> Schreb.	.	.	.	.	.	1	.	.	+	1	.	+	.	.	.	.	.	.	II ⁺¹
		<i>Alyssum alyssoides</i> (L.) L.	.	.	.	.	.	.	.	+	+	.	.	.	.	.	.	.	.	.	I ⁺
Autres taxons																					
		<i>Trisetum flavescens</i> (L.) P.Beauv.	.	.	.	.	.	.	.	.	.	+	.	+	.	.	.	.	.	.	I ⁺
		<i>Helictochloa pratensis</i> (L.) Romero Zarco	.	.	.	.	.	.	.	.	.	.	+	.	.	.	.	.	.	.	I ⁺
		<i>Pimpinella saxifraga</i> L.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	+	.	I ⁺
C		<i>Potentilla reptans</i> L.	.	.	.	.	.	.	.	.	+	.	.	.	.	.	.	.	.	.	I ⁺
		<i>Avenula pubescens</i> (Huds.) Dumort.	.	.	.	.	.	.	.	.	.	.	.	.	+	.	.	.	.	.	I ⁺
		<i>Plantago lanceolata</i> L.	.	.	.	.	.	.	.	.	.	.	+	.	+	.	.	+	.	.	I ⁺
		<i>Achillea millefolium</i> L.	.	.	.	.	.	.	+	.	.	.	.	.	.	.	.	.	.	.	I ⁺
		<i>Dactylis glomerata</i> L.	.	.	.	.	.	.	.	.	+	.	+	.	+	.	.	.	.	.	I ⁺
		<i>Bromus hordeaceus</i> L.	.	.	.	.	.	.	.	.	+	.	1	.	1	.	.	.	.	.	I ⁺¹
		<i>Taraxacum</i> sect. <i>Taraxacum</i>	.	.	.	.	.	.	.	.	.	.	+	.	+	.	.	.	.	.	I ⁺
		<i>Medicago sativa</i> L.	.	.	.	.	.	.	.	.	+	.	.	+	.	.	.	.	1	.	I ⁺¹
F		<i>Arrhenatherum elatius</i> (L.) P.Beauv. ex J.Presl & C.Presl	.	.	.	.	.	.	.	.	.	.	.	.	.	+	.	.	.	.	I ⁺
		<i>Cerastium fontanum</i> Baumg.	.	.	.	.	.	.	.	.	.	.	+	.	.	.	.	.	.	.	I ⁺
		<i>Trifolium pratense</i> L.	+	.	.	.	.	.	.	.	.	.	.	.	.	.	+	.	.	.	I ⁺
		<i>Crepis foetida</i> L.	.	.	.	.	.	.	.	.	.	.	.	+	.	+	.	.	.	.	I ⁺
		<i>Euphorbia flavicoma</i> DC.	.	+	.	.	1	.	.	.	.	.	.	.	.	.	.	.	.	.	I ⁺¹
		<i>Carlina acanthifolia</i> All.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	+	1	.	.	I ⁺¹
		<i>Anacamptis pyramidalis</i> (L.) Rich.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	+	.	.	I ⁺
		<i>Xanthoselinum alsaticum</i> (L.) Schur	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	+	I ⁺
		<i>Ophrys aranifera</i> Huds.	.	.	.	+	+	.	.	.	.	.	.	.	.	.	.	+	.	.	I ⁺
		<i>Xeranthemum cylindraceum</i> Sm.	.	.	.	.	.	.	.	+	.	.	.	.	.	.	.	.	.	.	I ⁺
		<i>Helleborus foetidus</i> L.	.	.	.	+	.	.	.	.	.	+	.	.	.	.	.	.	.	.	I ⁺
		<i>Salvia verbenaca</i> L.	.	.	.	.	.	.	.	.	.	.	1,2	.	1	.	.	.	.	.	I ¹
		<i>Tragopogon crocifolius</i> L.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	+	.	.	I ⁺
		<i>Stachys heraclea</i> All.	.	.	.	.	.	.	.	.	.	.	.	.	1	.	.	.	.	.	I ¹
		<i>Vicia tenuifolia</i> Roth	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	1	.	+	I ⁺¹
		<i>Bupleurum falcatum</i> L.	.	.	.	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	I ¹
		<i>Clinopodium acinos</i> (L.) Kuntze	+	.	.	.	+	.	.	.	.	.	.	.	.	.	.	.	.	.	I ⁺
		<i>Neottia ovata</i> (L.) Bluff & Fingerh.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	+	.	.	I ⁺
		<i>Orchis militaris</i> L.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	+	.	.	I ⁺
		<i>Silene otites</i> (L.) Wibel	.	.	.	.	.	.	+	.	.	+	.	.	.	.	.	.	.	.	I ⁺
		<i>Melampyrum arvense</i> L.	.	.	.	.	.	.	.	.	.	.	+	.	.	.	.	.	.	.	I ⁺
		<i>Falcaria vulgaris</i> Bernh.	.	+	+	.	.	.	.	.	.	.	.	.	.	.	.	.	1	.	I ⁺¹
		<i>Galatella linosyris</i> (L.) Rchb.f.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	+	.	.	.	I ⁺
		<i>Ophrys insectifera</i> L.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	+	.	.	I ⁺
		<i>Reseda lutea</i> L.	.	.	.	.	.	.	.	+	.	.	.	.	.	.	.	.	.	.	I ⁺
		<i>Bombycilaena erecta</i> (L.) Smoljan.	.	.	.	.	.	.	.	.	+	.	.	.	.	.	.	.	.	.	I ⁺
		<i>Veronica arvensis</i> L.	.	.	.	.	.	.	+	.	.	+	+	.	.	.	.	.	.	.	I ⁺
		<i>Cerastium pumilum</i> Curtis	.	.	.	.	.	.	.	+	.	+	.	.	.	.	.	.	.	.	I ⁺
		<i>Polygala</i> L.	.	.	.	1	.	.	.	.	.	.	.	.	.	.	+	.	.	.	I ⁺¹
		<i>Echium vulgare</i> L.	.	.	.	.	.	.	+	.	.	.	.	.	.	.	.	.	.	.	I ⁺